

Effects of dietary macronutrients on liver fat content in adults

Citation for published version (APA):

Winters-van Eekelen, E., Verkouter, I., Peters, H. P. F., Alssema, M., de Roos, B. G., Schrauwen-Hinderling, V. B., Roumans, K. H. M., Schoones, J. W., Zock, P. L., Schrauwen, P., Rosendaal, F. R., Dekkers, O. M., & de Mutsert, R. (2021). Effects of dietary macronutrients on liver fat content in adults: a systematic review and meta-analysis of randomized controlled trials. *European Journal of Clinical Nutrition*, 75(4), 588-601. <https://doi.org/10.1038/s41430-020-00778-1>

Document status and date:

Published: 01/04/2021

DOI:

[10.1038/s41430-020-00778-1](https://doi.org/10.1038/s41430-020-00778-1)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Effects of dietary macronutrients on liver fat content in adults: a systematic review and meta-analysis of randomized controlled trials

Esther Winters-van Eekelen¹ · Inge Verkouter¹ · Harry P. F. Peters² · Marjan Alssema² · Babette G. de Roos¹ · Vera B. Schrauwen-Hinderling³ · Kay H. M. Roumans^{1,4} · Jan W. Schoones⁵ · Peter L. Zock² · Patrick Schrauwen⁴ · Frits R. Rosendaal¹ · Olaf M. Dekkers^{1,6,7} · Renée de Mutsert¹

Received: 19 December 2019 / Revised: 3 September 2020 / Accepted: 2 October 2020
© The Author(s), under exclusive licence to Springer Nature Limited 2020

Abstract

Dietary macronutrient composition may affect hepatic liver content and its associated diseases, but the results from human intervention trials have been equivocal or underpowered. We aimed to assess the effects of dietary macronutrient composition on liver fat content by conducting a systematic review and meta-analysis of randomized controlled trials in adults. Four databases (PubMed, Embase, Web of Science, and COCHRANE Library) were systematically searched for trials with isocaloric diets evaluating the effect of dietary macronutrient composition (energy percentages of fat, carbohydrates, and protein, and their specific types) on liver fat content as assessed by magnetic resonance techniques, computed tomography or liver biopsy. Data on change in liver fat content were pooled by random or fixed-effects meta-analyses and expressed as standardized mean difference (SMD). We included 26 randomized controlled trials providing data for 32 comparisons on dietary macronutrient composition. Replacing dietary fat with carbohydrates did not result in changes in liver fat (12 comparisons, SMD 0.01 (95% CI -0.36; 0.37)). Unsaturated fat as compared with saturated fat reduced liver fat content (4 comparisons, SMD -0.80 (95% CI -1.09; -0.51)). Replacing carbohydrates with protein reduced liver fat content (5 comparisons, SMD -0.33 (95% CI -0.54; -0.12)). Our meta-analyses showed that replacing carbohydrates with total fat on liver fat content was not effective, while replacing carbohydrates with proteins and saturated fat with unsaturated fat was. More well-performed and well-described studies on the effect of types of carbohydrates and proteins on liver fat content are needed, especially studies comparing proteins with fats.

Introduction

Non-alcoholic fatty liver (NAFL) is clinically defined as a liver fat content of more than 5.6%, not due to excessive

alcohol consumption [1]. It is a major cause of chronic liver disease worldwide, associated with an increased risk of liver- and cardiovascular disease-related mortality [2–5]. Moreover, obesity and other features of the metabolic syndrome such as dyslipidaemia, insulin resistance and diabetes mellitus, are associated with NAFL [6–10]. The prevalence of NAFL continues to rise [2, 3] and has been estimated at 25% in adults [2], and between 65 and 85% in adults with obesity [11].

Supplementary information The online version of this article (<https://doi.org/10.1038/s41430-020-00778-1>) contains supplementary material, which is available to authorized users.

✉ Esther Winters-van Eekelen
E.van_eekelen@lumc.nl

¹ Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands

² Unilever Research and Development, Vlaardingen, the Netherlands

³ Department of Radiology, NUTRIM School for Nutrition, Toxicology and Metabolism, Maastricht University Medical Center, 6200 Maastricht, the Netherlands

⁴ Department of Nutrition and Movement Sciences, NUTRIM School for Nutrition and Translational Research in Metabolism, Maastricht University Medical Center, Maastricht, the Netherlands

⁵ Walaeus Library, Leiden University Medical Center, Leiden, the Netherlands

⁶ Department of Endocrinology, Leiden University Center, Leiden, the Netherlands

⁷ Department of Clinical Epidemiology, Aarhus University Hospital, Aarhus, Denmark

Since NAFL is still reversible, adequate treatment is needed to prevent the development into more severe forms of hepatic fat storage such as non-alcoholic steatohepatitis (NASH) [12, 13]. Drug-based treatments are primarily recommended for patients with a later stage of NAFL, whereas lifestyle changes are a cornerstone in guidelines on treatment of NAFL, including weight loss, eating healthier, and increasing physical exercise [12]. To date, interventions on NAFL mainly focus on decreasing total body fat by recommending calorie-restricted diets in overweight or obese patients [14–16]. However, besides diet quantity in the form of caloric restriction, macronutrient composition may be of importance, although evidence on this is scarce. Recent meta-analyses have shown that supplementation of omega-3 polyunsaturated fatty acids (PUFAs) is an effective intervention for reducing NAFL [17, 18].

Besides specific types of macronutrient such as omega-3 polyunsaturated fatty acids and fructose consumption, there are no meta-analyses on other macronutrients and other macronutrient types. In only one review on the effects of macronutrients on liver fat it has been described that a relatively high consumption of saturated fat increases the percentage of liver fat, whereas an increased consumption of refined sugars had no influence on liver fat [19]. However, the search of this review was limited and was not substantiated by a meta-analysis. Therefore, it remains unclear whether dietary macronutrients and their composition affect liver fat content. We aimed to assess the effect of dietary macronutrient composition on liver fat content, as measured by magnetic resonance imaging, proton magnetic resonance spectroscopy, computed tomography or liver biopsy, by performing a systematic review and meta-analysis of isocaloric randomized controlled trials in adults.

Methods

This systematic review and meta-analysis on dietary macronutrient composition and liver fat content was reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA-) guidelines and the recommendations of the Cochrane Collaboration [20, 21]. The protocol is registered at PROSPERO with registry ID number 100356.

Eligibility criteria

Databases were systematically searched for eligible publications based on a priori determined eligibility criteria. We systematically searched for randomized controlled dietary intervention trials evaluating the effect of macronutrient composition on liver fat content in adults. Liver fat was required to be reported as an outcome in title or abstract.

Studies including healthy adults as well as patients with obesity, metabolic syndrome, (pre)diabetes, NAFL or NASH and/or cardiovascular disease, were considered eligible. Trials that included individuals with malignant diseases or with alcoholic, drug-induced, viral or genetic causes of liver injury, were excluded.

Both macronutrient comparisons (carbohydrates versus fat, carbohydrates versus protein, protein versus fat) and macronutrient types comparisons (types of fat, types of carbohydrates, and types of protein) were assessed. Since several reviews and meta-analyses on omega-3 fatty acids and fructose have been published recently [17, 22–26], studies were excluded when the dietary intervention was primarily focused on these types of macronutrient comparisons. Studies that used hyper- or hypo-caloric interventions were only eligible when caloric intake was equal in both study arms. Furthermore, the interventions had to be provided for at least one week, since seven days of dietary intervention was deemed necessary to influence fat oxidation in the liver [27]. In addition, trials that involved co-interventions, such as exercise or other lifestyle interventions, were only included when similar in both arms of the trial. Trials solely providing their participants with dietary advice rather than food items, as well as trials presenting insufficient information on macronutrient composition were not eligible. Assessment methods of liver fat content were predefined: only trials in which liver fat content was measured by magnetic resonance imaging (MRI), magnetic resonance spectroscopy (MRS), computed tomography (CT) or liver biopsy were considered [28, 29]. A detailed overview of all eligibility criteria can be found in Supplementary Table 1.

Search strategy

We conducted a systematic search to identify eligible publications. In cooperation with a trained librarian (JWS), a detailed search strategy was composed for the four bibliographic databases: PubMed, Embase (OVID-version), Web of Science, and COCHRANE Library. The search query consisted of a combination of the following concepts: macronutrients (exposure terms), liver fat (outcome terms) and (randomized controlled) trials. The search strategy was adjusted for all consulted databases, taking into account the differences of the various controlled vocabularies as well the differences of database-specific technical variations (e.g., the use of quotation marks). Case reports, animal-only studies and conference abstracts were excluded. No restrictions were made on language and publication year. The final search was performed on February 19th, 2018 and repeated on June 17, 2019 and March 6, 2020. All search strings used can be found in the supplementary data.

Study selection process

First, duplicate publications were removed. Titles and abstracts of remaining identified publications were screened for eligibility by 6 reviewers (BdR, EW-vE, HP, IV, KR, MA) in preassembled pairs. Each reviewer of a pair independently screened and coded an assigned part of the articles ‘include’, ‘unclear’ or ‘exclude’. Disagreements on inclusion were discussed in the pre-assembled pairs until consensus was reached. Subsequently, potentially relevant publications were independently assessed in full-text by three reviewers (BdR, IV, EW-vE). In case of multiple publications of a single trial, the first published version was included. Discrepancies on the eligibility of articles were resolved by discussion until consensus was reached. The selection of publications was managed by the Rayyan QCRI web application (Qatura Computing Research Institute, 2016) [30].

Data collection and extraction

Data extraction was independently performed by two reviewers (EW-vE and IV) using a predefined sheet in Microsoft Excel, Version 15.40. Extracted data were compared and discrepancies were resolved. Data were extracted on four categories following the recommendations of the Cochrane Collaboration; characteristics of the study (i.e., dietary comparison, location, design), the participants (i.e., number of randomized/analyzed participants, sex, mean age, mean body weight, mean BMI), the dietary interventions (i.e., compositions, follow-up time) and the outcomes per arm of the trial [21].

Risk of bias assessment

Two reviewers (EW-vE and IV) independently assessed the risk of bias for included studies, using the Cochrane ‘Risk of bias’ tool for randomized controlled trials [24]. This tool involved a classification of six different domains of bias (i.e., selection bias, performance bias, attrition bias, detection bias, reporting bias, and (design-specific) other sources of bias) with seven corresponding domains: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting and “other sources of bias”. For detection of the “other sources of bias”, reviewers were in particular alert to (self) reporting bias, compliance assessment and carry-over effects in cross-over trials, with trials lacking a wash-out period being at higher risk. Each domain was separately judged as having a “low”, “high” or “unclear” risk of bias. In addition, a support for judgment was given and summarized following the criteria outlined by the Cochrane

Collaboration [21]. Any discrepancies in bias coding were resolved by discussion.

Direct pairwise meta-analyses

To perform meta-analyses for continuous outcomes measured with different measuring instruments of liver fat on different scales (i.e., MRS/MRI (%) and CT-scans (Hounsfield Units)), effect estimates were expressed as standardized mean difference (SMD) with corresponding 95% confidence interval (95% CI). When studies only reported relative changes in liver fat, the absolute change based on the relative change and the baseline value was calculated. If trials presented medians and interquartile ranges (IQRs), values were converted into means and standard deviations according to the Cochrane Collaboration [21].

Intervention effects were pooled by performing standard pairwise meta-analyses for all comparisons that contained at least three comparisons between diets. A random-effects model was used (method of DerSimonian and Laird [31]) for the comparison between a low-carbohydrate high-fat and a high-carbohydrate low-fat diet and due to the limited number of included studies a fixed-effect model for the other two comparisons. For the study of Luukkonen et al. [32], two interventions (saturated fat and unsaturated fat) were compared against the same control group (carbohydrates). To correct for these multiple correlated comparisons the number of participants in the control arm was divided by the number of comparisons (i.e., two) thereby creating two (reasonably independent) comparisons (Cochrane handbook Chapter 16.5.4). We performed a sensitivity analysis in which the two groups with physical activity as a co-intervention from the study of Bozzetto et al. were excluded to eliminate the potential effect of physical activity on the results. The diet that was expected to be beneficial, as described in the rationale of the included studies, was considered as the intervention arm (high unsaturated fat-low saturated fat, high protein-low carbohydrates and high-carbohydrates low-fat), and the other the control arm (saturated fat, high carbohydrates, and high fat). As a result, a negative standardized mean difference can be interpreted as a decrease in liver fat in the intervention arm compared with the control arm, which means that the intervention arm is favored. In case of an overfeeding design, a negative standardized mean difference represents a smaller increase in liver fat in the intervention arm compared to the control arm. A positive standardized mean difference indicates that the control arm is favored. Guidelines state that an SMD of 0.2 can be considered small, 0.5 as medium and 0.8 as high [33].

Statistical heterogeneity was assessed using the I-squared statistic [34, 35]. Heterogeneity was considered to be low if the I^2 value was under 40%, moderate if between 30 and

60%, substantial if between 50 and 90% and considerable when between 75 and 100% [24, 36]. All statistical analyses were conducted using Stata statistical Software (Statacorp, College Station, Texas, USA) version 14.

Handling missing data

In case of unreported or incomplete data on mean changes (or SD) in liver fat content between baseline and follow-up, the original investigators were contacted and asked to provide missing data. When no response was received, we calculated mean differences using standard deviations based on the information that was provided (baseline or follow-up value with corresponding SD), as described in a previous meta-analysis [37]. Trials were not included when relevant data to calculate mean differences was not provided [21].

Small-study effects

A funnel plot was used for graphical examination of small-study effects [38–41]. In addition, Egger's test was performed [24, 39, 41] if more than 10 studies for a specific analysis were available [42, 43].

Results

Study selection

Of the 4731 publications retrieved, a total of 3721 unique publications were screened on title and abstract (Fig. 1). Of those, 3594 publications were excluded after screening of titles and abstracts for eligibility. A total of 127 articles were assessed for eligibility based on full text, of which 101 were excluded due to the following reasons: no dietary intervention ($n = 25$), interventions not isocaloric ($n = 10$), multiple publications from a single trial ($n = 4$), no original research paper ($n = 17$), co-interventions not equal in both arms ($n = 2$), no adequate comparison ($n = 3$), no MRI/MRS/CT/biopsy liver fat outcome ($n = 28$), population younger than 18 years ($n = 4$) or no RCT design ($n = 8$), leaving a total of 26 included articles [32, 34, 38, 39, 42, 44–64] (Fig. 1).

For one study, only two out of three arms were incorporated into the meta-analysis, as the diet in one arm contained less calories than the diet in the other two arms [54]. Ultimately, 32 eligible comparisons remained for analyses as four studies contained more than one comparison [32, 38, 46, 64].

Study characteristics

Table 1 shows the characteristics of the 26 randomized controlled trials. Studies were published between 2002 and

2019 and the number of participants ranged from 7 to 166. The duration of the studies varied between 7 days and two years. With regard to the macronutrient comparisons, ten studies reported effects of a low-carbohydrate high-fat (LCHF, En% carbohydrates ranging from 10 to 40, fat 42 to 75)-diet compared with a high-carbohydrate low-fat (HCLF, En% carbohydrates ranging from 53 to 65, fat 16 to 34)-diet [32, 38, 39, 42, 46–48, 56–58] (Supplementary Table 2). Five studies compared a low-protein high-carbohydrate (LPHC, protein 5–18.5 and carbohydrates 45–60 En%)-diet with a high-protein low-carbohydrate (HPLC, protein 22.1–30.5 and carbohydrates 29.7–41 En%)-diet [50, 53, 54, 61, 63]. There were no studies on the comparisons between fat and protein content of the diet.

The other studies performed comparisons between types of macronutrients. A total of six studies compared different types of dietary fat, of which four studies compared a diet high in saturated fatty acids (SFAs) with a diet high in unsaturated fatty acids (UFAs) [32, 34, 55, 62], one study compared trans fatty acids with palm- and sunflower oil [45] and one study looked at replacement of long-chain fatty acids with medium-chain fatty acids [52]. In two studies dietary fibers were compared with other carbohydrates [38, 44], one study compared whole grain wheats with refined wheats [59], another compared milk with yogurt [60], one compared increased fiber intake, a diet without red meat and a control diet [64], and in two studies diets containing animal protein was compared with diets containing plant/soy protein [49, 51].

In total, twenty studies used a parallel design, whereas six had a cross-over design [44, 48, 51, 54, 57, 63]. Two studies assessed the liver fat content using CT [52, 53], whereas all other studies used MRS/MRI. One study assessed liver fat content both with MRI and MRS, of which we chose to use the MRS results in the meta-analysis as this is considered the most reliable method [11]. Most studies mainly included participants with overweight or obesity, varying from adolescents to elderly, except for six studies that included lean participants [44, 48, 50, 52, 54, 55] (Table 1). Additional information on the macronutrient composition per study arm can be found in Supplementary Table 2.

Risk of bias

The risk of bias assessment for included studies can be found in Table 2. In eleven studies there was high risk of performance bias, in two studies there was high risk of detection bias, in four studies of attrition bias, in seven studies of reporting bias and in nine studies there was a high risk of other bias.

The majority of the studies had an unclear risk of selection bias due to a lack of information on concealment




Fig. 1 Flowchart of included randomized controlled trials in meta-analysis on dietary macronutrient composition in relation to liver fat. The flow diagram displays the number of records identified, and the different phases of inclusion and exclusion with the corresponding reasons.

of allocation. Overall, there was unclear risk of selection bias and detection bias, and substantial risk of performance, attrition, reporting and other types of bias.

Effects of interventions

Table 3 provides a summary of findings for all included trials. It also shows the changes in liver fat content and

corresponding SMDs for all studies individually. Based on all included trials, we were able to perform three meta-analyses, as described below. A total of 21 studies were included, comprising a total of 25 comparisons between different diets. As we decided to only perform a meta-analysis on exchanges that contained at least three comparisons between dietary intervention arms, we could not meta-analyze comparisons of trans fats with palm- and

Table 1 Characteristics of randomized controlled trials included in meta-analysis on association between dietary macronutrient composition and hepatic triglyceride content.

Author, year	Study design	Length (days)	Run-in/ wash-out	Liver fat measurement	Men (%) or mean age (y)	BMI range or mean at baseline (kg/m ²)	Intervention	N	Control	N	
Bawden, 2016	Cross-over	7	No/Yes	¹ H-MRS	100	20.1	23.0	LGI (high fiber)	7	HGI (low fiber)	7
Bendsen, 2011	Parallel	112	No/NA	¹ H-MRS	0	45–70	25–32	Trans fatty acids	23	Palm/sunflower oil	23
Bjermo, 2012	Parallel	70	No/NA	¹ H-MRS	34	30–65	30.8 ^a	Polysaturated fat	28	Saturated fat	28
Bozzetto, 2012a	Parallel	56	Yes/NA	¹ H-MRS	75.0	35–70	29.1 ^a	Carbohydrates/fiber	9	Monounsaturated fat	8
Bozzetto, 2012b							30.5 ^a	Carbohydrates/fiber (+ exercise)	10	Monounsaturated fat (+ exercise)	9
Chen, 2019	Parallel	168	No/NA	¹ H-MRS	0	36–66	>28	Yogurt	20	Milk	20
Errazuriz, 2017	Parallel	84	Yes/NA	¹ H-MRS	53.3	61.7	31.7 ^a	Control/carbohydrates	11	Monounsaturated fat	15
Gepner, 2019	Parallel	~180	No/NA	MRI	85.4	47.7	30.9 ^a	Fiber	13	Control	11
Haufe, 2017	Parallel	~180	No/NA	¹ H-MRS	17.6		>25	Low fat/high carbohydrates	50	High fat/low carbohydrates	52
Herpen, 2011	Parallel	42	Yes/NA	¹ H-MRS	100	55.2	28.8 ^a	Low fat/high carbohydrates	9	High fat/low carbohydrates	9
Kirk, 2009	Parallel	42	No/NA	¹ H-MRS	18.2	43.6	36.5	Low fat/high carbohydrates	11	High fat/low carbohydrates	11
Luukkonen, 2018	Parallel	21	No/NA	¹ H-MRS	44.7	48.0	31.0	Carbohydrates	12	Unsaturated fat	12
Luukkonen, 2018								Carbohydrates	12	Saturated fat	14
Luukkonen, 2018								Unsaturated fat	12	Saturated fat	14
Marin-Alejandre, 2019	Parallel	~180	No/NA	MRI	52	50	33.5 ^a	FLiO diet (high protein)	39	AHA diet (high carbohydrates)	37
Marina, 2014	Cross-over	28	Yes/Yes	¹ H-MRS	76.9	36.0	33.6	Low fat/high carbohydrates	10	High fat/low carbohydrates	10
Markova, 2017	Parallel	42	No/NA	¹ H-MRS	64.9	49–78	30.2 ^a	Plant protein	19	Animal protein	18
Martens, 2014	Parallel	84	No/NA	¹ H-MRS	33.3	24.0	22.9	High protein/low carbohydrates	7	Low protein/high carbohydrates	9
Nosaka, 2002	Parallel	28	No/NA	CT	100	27–51	23.1 ^a	Long-chain triacylglycerols	11	Medium-chain triacylglycerol	11
Ooi, 2015	Parallel	~730	No/NA	CT	0	70–80	26.5 ^a	Protein	82	Carbohydrates	84
Rietman, 2014	Cross-over	14	Yes/No	¹ H-MRS	70.4	22.8	21.5	High protein/low carbohydrates	17	Normal protein/normal carbohydrates	17
Rosqvist, 2014	Parallel	49	No/NA	MRI	70.3	20–38	20.3 ^a	Polysaturated fat	18	Saturated fat	19
Rosqvist, 2019	Parallel	56	No/NA	MRI	61.7	42	28.0 ^a	Polysaturated fat	30	Saturated fat	30
Schutte, 2018	Parallel	84	Yes/NA	¹ H-MRS	62.0	45–70	27.81	Whole grain wheat	20	Refined wheats	18

Table 1 (continued)

Author, year	Study design	Length (days)	Run-in/ wash-out	Liver fat measurement	Men (%) or mean age (y)	BMI range or mean at baseline (kg/m ²)	Intervention	N	Control	N	
Skyyte, 2019	Cross-over	42	No/No	¹ H-MRS	71.4	64	30.1	High protein	27	Control (high carbohydrates)	27
Utzschneider, 2013	Parallel	28	No/NA	¹ H-MRS	37.0	69.3	27.4 ^a	Low fat/LGI	20	High fat/HGI	15
Van Niel, 2014	Cross-over	28	Yes/NA	¹ H-MRS	0	61.0		Soy protein	10	Mixed protein	10
Westerbacka, 2005	Cross-over	14	Yes/ Unclear	¹ H-MRS	0	43	33.0	Low fat/high carbohydrates	10	High fat/low carbohydrates	10
Willmann, 2019	Parallel	~180	No/NA	¹ H-MRS	34.8	42	31.2	Fiber	44	Control	37
Willmann, 2019								No red meat	41	Control	37
Willmann, 2019								Fiber	44	No red meat	41

^aWeighted mean BMI based on mean baseline BMI values of separate arms.

BMI body mass index, *CHO* carbohydrates, *CT* computed tomography, ¹*H*-*MRS* proton magnetic resonance spectroscopy, *HGI* high glycaemic index, *LGI* low glycaemic index, *MRI* magnetic resonance imaging, *MUFA* monounsaturated fatty acids, *PUFA* polyunsaturated fatty acids, *SFA* saturated fatty acids.

sunflower oil, long chain with medium-chain fat, dietary fiber with other carbohydrates, whole grain wheats with refined wheats, and animal protein with plant protein. Due to the limited number of included trials, we were not able to perform subgroup analyses on disease state, sex, ethnicity or study duration. Moreover, as there were no studies comparing dietary protein with fat, we could not perform a network meta-analysis in which all macronutrients could be compared both directly and indirectly [38, 44, 45, 49, 51, 52].

High-carbohydrate low-fat versus low-carbohydrate high-fat diets

Out of 12 comparisons for a low-carbohydrate high-fat with a high-carbohydrate low-fat diet, three comparisons favored a low-carbohydrate high-fat diet over a high-carbohydrate low-fat diet [38, 46], while two other comparisons showed the opposite [42, 57] (Fig. 2). The other studies showed no difference. Heterogeneity was substantial (67.8%). No small study effects seemed to be present (Supplementary Fig. 1) (*P* value for Egger's test 0.58). The overall pooled effect of high-carbohydrate low-fat versus high-fat low-carbohydrate was: SMD 0.01, 95% CI -0.36; 0.37 (Fig. 2).

After excluding the two groups with a co-intervention of physical exercise from the study of Bozzetto, results were similar (data not shown).

Dietary saturated fat versus unsaturated fat

Only three studies examined the effect of unsaturated fat compared to saturated fat, of which all three found that an unsaturated fat diet reduces liver fat compared with saturated fat [32, 34, 55] (Fig. 3). The overall effect showed that unsaturated fat as compared with saturated fat reduced liver fat to a large extent (SMD -0.75, 95% CI -1.11; -0.39. A funnel plot is shown in Supplementary Fig. 2; Egger's test was not performed due to an insufficient number of included studies.

High-protein low-carbohydrate versus low-protein high-carbohydrate diets

Three studies assessed the effect of a high protein-low carbohydrate compared to a low-protein high-carbohydrate diet on liver fat. One study found that a high-protein low-carbohydrate diet resulted in reduced liver fat content compared to a low-protein high-carbohydrate diet [50], whereas the other two studies did not find a difference [53, 54] (Fig. 4). The overall pooled effect showed that a high-protein low-carbohydrate diet moderately reduced liver fat as compared to a low-protein high-carbohydrate

Table 2 Risk of bias of randomized controlled trials included in systematic review and meta-analysis on macronutrient and macronutrient types composition in relation to liver fat content in adults of 18 years and older.

First author	Selection bias		Performance bias	Detection bias	Attrition bias	Reporting bias	Other bias
	Random sequence generation	Allocation concealment					
Bawden, 2017	Unclear	Unclear	Unclear	Low	Unclear	Unclear	Low
Bendsen, 2011	Low	Low	Low	Low	Low	Unclear	Unclear
Bjermo, 2012	Unclear	Unclear	High	Low	High	Unclear	High
Bozzetto, 2012	Low	Low	Unclear	Low	Low	Unclear	High
Chen, 2019	Low	Unclear	High	Low	Low	Low	High
Errazuriz, 2007	Unclear	Unclear	Unclear	Low	Low	High	Low
Gepner, 2019	Unclear	Unclear	High	Low	High	Unclear	High
Haufe, 2017	Low	Low	High	Unclear	High	Unclear	Unclear
Herpen, 2011	Low	Unclear	Unclear	Unclear	Low	High	High
Kirk, 2009	Unclear	Unclear	Unclear	Unclear	Low	Unclear	High
Luukkonen, 2018	Unclear	Unclear	High	High	Low	Low	Low
Marin-Alejandre, 2019	Unclear	Unclear	High	Unclear	Unclear	Low	Unclear
Marina, 2014	Unclear	Unclear	Unclear	Low	High	High	High
Markova, 2017	Low	Unclear	High	High	Unclear	Low	Unclear
Martens, 2014	Low	Low	High	Unclear	High	High	Low
Nosaka, 2002	Low	Unclear	Low	Unclear	Low	Unclear	Unclear
Ooi, 2015	Low	Low	Low	Low	Low	High	Low
Rietman, 2014	Low	Low	Low	Unclear	Low	High	Unclear
Rosqvist, 2014	Low	Unclear	Low	Low	Unclear	Unclear	Unclear
Rosqvist, 2019	Low	Low	Low	Low	Unclear	Unclear	Unclear
Schutte, 2018	Unclear	Unclear	Low	Unclear	High	Unclear	Low
Skytte, 2019	High	Unclear	High	Low	Unclear	Unclear	High
Utzschneider, 2012	Unclear	Unclear	Low	Low	Low	Unclear	Unclear
van Niel, 2014	Unclear	Unclear	High	Low	Unclear	Low	Low
Westerbacka, 2005	Low	Unclear	Unclear	Low	Low	High	High
Willmann, 2019	Low	Low	High	Unclear	High	Low	Unclear

diet (SMD -0.32 , 95% CI -0.58 ; -0.05). A funnel plot is shown in Supplementary Fig. 3, Egger's test was not performed due to an insufficient number of included studies.

Discussion

With this systematic review and meta-analysis including randomized controlled trials we have provided a summary of the evidence on the effect of dietary macronutrient composition on the amount of liver fat, as assessed by ^1H -MRS, MRI or CT. Our results show that replacing dietary fat with carbohydrates did not result in changes in liver fat. Diets high in unsaturated fat lead to a larger decrease (or smaller increase in case of an overfeeding design) in liver fat content than diets high in saturated fat. A high-protein low-carbohydrate diet reduces liver fat as compared with a low-protein high-carbohydrate diet.

Our results focusing on liver fat content are in line with the review of Parry and Hodson, in which the authors describe that most studies suggest no influence on liver fat by diets that are high in carbohydrates in the form of free sugars [19]. The increase in liver fat observed in diets high in fat seems to be attributable to an increased saturated fat consumption, while increased consumption of mono- or polyunsaturated fat may reduce liver fat content [19], which supports the results of our meta-analysis. The beneficial effects of unsaturated fat on liver fat content compared to saturated fat were also reported in another recent review [65]. Additionally, results from this meta-analysis are in agreement with the findings from a meta-analysis on the effects of mutual exchanges of different dietary fats and carbohydrates on glucose-insulin homeostasis, an outcome strongly related to NAFL. The authors of this meta-analysis found that replacement of carbohydrates or saturated fat with polyunsaturated fat led to an improved insulin

Table 3 Standardized mean differences of randomized controlled trials included in meta-analysis on association between dietary macronutrient composition and hepatic triglyceride content.

Author	Intervention	N	Change in liver fat after intervention (% or HU)	Control	N	Change in liver fat after intervention (% or HU)	Mean difference in change in liver fat between arms (intervention-control, % or HU)	Standard deviation of mean difference	Standardized mean difference
Bawden, 2016	Fibre	7	-0.4	Other carbs	7	1.3	-1.70	1.46	-1.16
Bendsen, 2011	Palm/sunflower oil	23	-0.6	Trans fatty acids	23	-0.8	0.20	4.10	0.05
Bjermo, 2012	PUFA	28	-0.9	SFA	28	0.3	-1.20	2.01	-0.60
Bozzetto, 2012a	Low fat	9	-1.6	Low carb	8	2.2	0.60	0.58	1.04
Bozzetto, 2012b	Low fat	10	0.1	Low carb	9	-2.5	2.60	2.70	0.96
Chen, 2019	Yogurt	20	-4.97	Milk	20	0.5	-5.47	7.1	-0.77
Errazuriz, 2017	Low fat	11	0.7	Low carb	15	-1.7	2.40	1.75	1.37
Errazuriz, 2017	Fibre	13	-0.6	Other carbs	11	0.7	-1.30	1.33	-0.98
Gepner, 2019	Low fat	79	-5.8	Low carb	78	-7.3	1.5	5.31	0.29
Haufe, 2017	Low fat	50	-4.0	Low carb	52	-3.6	-0.40	4.31	-0.09
Herpen, 2011	Low fat	9	-0.52	Low carb	9	0.37	-0.89	0.88	-1.01
Kirk, 2009	Low fat	11	-4.98	Low carb	10	-4.71	-0.27	1.35	-0.20
Luukkonen, 2018	UFA	12	0.79	SFA	14	2.72	-1.93	1.76	-1.10
Luukkonen, 2018	Low fat	12	1.37	Low carb	14	2.72	-1.35	1.77	-0.76
Luukkonen, 2018	Low fat	12	1.37	Low carb	12	0.79	0.58	1.78	0.32
Marin-Alejandre, 2019	High protein	39	-4.2	High carb	37	-3.6	-0.6	6.9	-0.09
Marina, 2014	Low fat	10	-2.2	Low carb	10	-1.3	-0.90	2.28	-0.39
Markova, 2017	Plant protein	17	-6.8	Animal protein	15	-6.7	-0.10	8.96	-0.01
Martens, 2014	High protein	7	-0.03	High carb	9	0.05	-0.08	0.08	-1.05
Nosaka, 2002	Long chain FA	11	0.03	Medium chain FA	11	0.02	0.01	0.10	0.1
Ooi, 2015	High protein	82	0.00	High carb	84	0.04	-0.04	0.20	-0.2
Rietman, 2014	High protein	17	-0.05	High carb	17	0.11	-0.16	0.26	-0.62
Rosqvist, 2014	PUFA	18	0.04	SFA	19	0.56	-0.52	0.71	-0.73
Rosqvist, 2019	PUFA	30	-0.09	SFA	30	1.54	-1.63	1.82	-0.90
Utzschneider, 2013	Low fat	20	-0.50	Low carb	15	0.4	-0.9	2.50	-0.36
Van Nielen, 2014	Soy protein	10	-0.4	Meat protein	10	-0.9	0.5	0.84	0.59
Schutte, 2018	Whole grain wheats	20	0.53	Refined grain wheats	18	2.00	-1.47	2.00	-0.72
Westerbacka, 2005	Low fat	10	-2.0	Low carb	10	3.5	-5.5	5.62	-0.98
Willmann, 2019	Fiber	44	-1.7	Control	37	-0.8	-0.9	10.4	-0.09
Willmann, 2019	No red meat	41	-2.1	Control	37	-0.8	-1.3	22.6	-0.06
Willmann, 2019	Fiber	44	-1.7	No red meat	41	-2.1	0.4	14.6	0.03

Standardized mean difference (SMD) was calculated by dividing the mean difference between the arms by the standardized deviation of the difference between the arms.

FA fatty acid, HU Hounsfield Unit, MUFA mono-unsaturated fatty acids, PUFA poly-unsaturated fatty acids, SFA saturated fatty acids.

secretion capacity, lower fasting glucose, improved Homeostatic Model Assessment for Insulin Resistance (HOMA-IR) and lower hemoglobin A1C (HbA1c) [65]. The exchange of saturated fat for carbohydrates did not

affect most outcomes, except for a decrease in fasting insulin [66].

Although the pathogenesis of liver fat accumulation is not completely elucidated yet, it is assumed that both high




Fig. 2 Difference between effects of a low-carbohydrate high-fat diet (LCHF) and a high-carbohydrate low-fat (HCLF) on liver fat content in studies included in meta-analysis: a random-effects model. Standardized mean difference (SMD) was calculated by dividing the mean difference between the arms by the standardized

deviation of the difference between the arms. A negative standardized mean difference can be interpreted as a decrease in liver fat in the intervention arm compared with the control arm, which means that the intervention arm is favored.




Fig. 3 Difference between effects of a diet high in saturated fats (SFA) and a diet high in unsaturated fat (UFA) on liver fat content in studies included in meta-analysis: a fixed-effects model. Standardized mean difference (SMD) was calculated by dividing the mean difference between the arms by the standardized deviation of the

difference between the arms. A negative standardized mean difference can be interpreted as a decrease in liver fat in the intervention arm compared with the control arm, which means that the intervention arm is favored.

caloric intake and dietary composition influence liver fat content. Dietary intake of specific nutrients (e.g., fructose) may increase de novo lipogenesis, and together with increased lipolysis of visceral fat this may contribute to an

increased flux of free fatty acids in the liver, leading to hepatic fat accumulation [10, 67]. Additionally, n-6 polyunsaturated fatty acids have been suggested to suppress lipogenic gene expression and could thereby decrease de




Fig. 4 Difference between effects of a low-protein high-carbohydrate (LPHC) diet and a high-protein low-carbohydrate (HPLC) diet on liver fat content in studies included in meta-analysis: a fixed-effects model. Standardized mean difference (SMD) was calculated by dividing the mean difference between the arms by the

standardized deviation of the difference between the arms. A negative standardized mean difference can be interpreted as a decrease in liver fat in the intervention arm compared with the control arm, which means that the intervention arm is favored.

novo lipogenesis and thereby decrease accumulation of liver fat [68], which is consistent with the findings of this meta-analysis showing that this holds true more generally for unsaturated fat and that exchanging saturated for unsaturated fat can lower liver fat.

A strength of this study is that it is the first comprehensive meta-analysis on the effect of macronutrient composition and macronutrient types on liver fat. The review process has been performed systematically and only studies in which liver fat was measured with either MRI, ¹H-MRS, or CT were included. Moreover, we only included studies that performed a dietary intervention rather than only providing dietary advice.

This study also has some limitations. The first one is that comparing and meta-analyzing data from different dietary intervention trials appeared challenging, as there was considerable heterogeneity in study duration and composition of the diets, percentages of macronutrients exchanged, and total amount of energy of provided diets (hyper-, hypo- or isocaloric), which might have attenuated the effects. Whereas some studies specified which subtypes of dietary fats or carbohydrates were replaced, others did not, making the interpretation of the results difficult. As our results on exchanging unsaturated with saturated fat have shown, the fat type that is replacing the carbohydrates is likely relevant. Three randomized trials [32, 38, 46] replaced carbohydrates with unsaturated fats and show that a low-carbohydrate high-fat diet leads to less liver fat compared with a high-carbohydrate low-fat diet, whereas most other studies suggest that a high-carbohydrate low-fat diet leads to less liver

fat. However, information on the type of dietary fat used to replace carbohydrates was lacking in most studies, as was information on quality and type of carbohydrates (complex, refined or fibers). Even in a neutral energy balance, de novo lipogenesis increased as a result of diets high in fructose as compared with complex carbohydrates diet [69]. Therefore, results could have been both under- and overestimated.

Moreover, this meta-analysis focused on the exchange between two macronutrient (subtypes) irrespective of the energy percentage derived from these specific macronutrients. Therefore, the studies show marked heterogeneity in the percentual energy contribution of the macronutrient subtypes that were exchanged. Studies with a larger exchanged energy percentage of macronutrients between the compared diets may have resulted in larger effect estimates than studies with smaller exchanges in energy percentages. However, the effect sizes of the studies were not proportional to the amount of energy percentage that was exchanged.

Additionally, total caloric intake varied considerably between studies. Whereas some studies used an overfeeding design in which participants were instructed to consume more calories than their usual diet, other studies used an isocaloric or hypocaloric diet. Our only criterion regarding energy intake was that it should be equal in both study arms within a trial, regardless of whether energy intake was below, above or equal to the energy requirement of the participants. Therefore, mean caloric intake varied from 1.100 kilocalories per day [47] to over 3.400 kilocalories per day [54]. Although the number of included arms was too small to perform stratified analyses, the effect of

macronutrient composition did not seem to be modified by caloric intake after visual inspection in the meta-analysis on dietary carbohydrates versus fat, which included the most comparisons.

A second limitation of this review is that data of variance within the dietary arms of the included trials (e.g., variance of mean change in liver fat or variance of mean difference) were not always reported. Therefore, P-values of the mean differences in change in liver fat – that were converted to corresponding *t*-values – had to be used to calculate the standard deviations, standard error of the means and the 95% CIs of the mean differences in change in liver fat by Cochrane equations [36]. With these calculated values, mean differences could be converted to standardized mean differences and their corresponding 95% CIs. However, some studies did not present exact P-values of the mean difference, but exclusively presented the level of significance (e.g., $P < 0.05$ or $P < 0.01$). As described by the Cochrane Handbook, the limits of the significance level were used for these trials as a conservative approach [36]. This approach may have caused imprecision of the variance for each trial, which is reflected in a larger confidence interval around the SMD and a decreased weight of the study [36].

As only a limited number of studies could be included in this meta-analysis, we recommend that more large randomized controlled dietary trials with a low risk of bias and of sufficient power are performed, in which complete and transparent reporting of results is of great importance in order to address this gap in knowledge. Especially trials in which proteins and fats are exchanged are warranted, as they were completely lacking. The sources and types of these macronutrients should be well-specified. Bridging this gap in research is essential for the development of preventive strategies for fatty liver in the future.

In conclusion, this systematic review and meta-analysis of randomized controlled trials showed that replacing total carbohydrates with total fats has no effect on liver fat content. Replacing saturated fat with unsaturated fat resulted in a decrease or a smaller increase in liver fat content, and replacing carbohydrates with proteins also seems to lead to less liver fat. Only a limited number of eligible studies could be included, which supports an essential need for additional experimental studies on dietary macronutrient composition and liver fat content in order to provide optimal prevention and treatment for non-alcoholic fatty liver by dietary interventions.

Data availability

Data used in the manuscript, code book, and analytic code will be made available to editors upon request.

Acknowledgements We thank P.K. Luukkonen, S. Haufe, and M.R. Mensink for providing additional data for our analyses. We acknowledge the support from the Netherlands Cardiovascular Research Initiative: an initiative with support of the Dutch Heart Foundation (CVON2014-02 ENERGISE). This research was in part financed by the Ministry of Economic Affairs and Climate Policy by means of the PPP Allowance made available by the Top Sector Life Sciences & Health to stimulate public-private partnerships and by Unilever R&D Vlaardingen.

Author contributions EW-vE, IV, RdM, MA, HP, PZ, FR, VS-H, and PS designed research, EW-vE, IV, BdR, RdM, HP, MA, JS and KR conducted research, EW-vE, IV, BdR, and OD analyzed data, EW-vE, IV, BdR, RdM, HP, and MA wrote paper, EW-vE and RdM had primary responsibility for final content. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest MA, HP, and PZ were employees of Unilever R&D Vlaardingen at the time of this study. The other authors declare no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Petäjä EM, Yki-Järvinen H. Definitions of normal liver fat and the association of insulin sensitivity with acquired and genetic NAFLD—a systematic review. *Int J Mol Sci.* 2016;17:633.
2. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease—meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology.* 2016;64:73–84.
3. Williams CD, Stengel J, Asike MI, Torres DM, Shaw J, Contreras M, et al. Prevalence of nonalcoholic fatty liver disease and non-alcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology.* 2011;140:124–31.
4. Haddad TM, Hamdeh S, Kanmanthareddy A, Alla VM. Non-alcoholic fatty liver disease and the risk of clinical cardiovascular events: a systematic review and meta-analysis. *Diabetes Metab Syndrome.* 2017;11:S209–16.
5. Targher G, Day CP, Bonora E. Risk of cardiovascular disease in patients with nonalcoholic fatty liver disease. *N Engl J Med.* 2010;363:1341–50.
6. Chalasani N, Younossi Z, Lavine JE, Diehl AM, Brunt EM, Cusi K, et al. The diagnosis and management of non-alcoholic fatty liver disease: Practice Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. *Hepatology.* 2012;55:2005–23.
7. Mantovani A, Byrne CD, Bonora E, Targher G. Nonalcoholic fatty liver disease and risk of incident type 2 diabetes: a meta-analysis. *Diabetes Care.* 2018;41:372–82.
8. Vanni E, Bugianesi E, Kotronen A, De Minicis S, Yki-Järvinen H, Svegliati-Baroni G. From the metabolic syndrome to NAFLD or vice versa? *Digestive Liver Dis.* 2010;42:320–30.
9. Papandreou D, Andreou E. Role of diet on non-alcoholic fatty liver disease: an updated narrative review. *World J Hepatol.* 2015;7:575.

10. Buzzetti E, Pinzani M, Tsochatzis EA. The multiple-hit pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Metabolism*. 2016;65:1038–48.
11. Fabbriani E, Sullivan S, Klein S. Obesity and nonalcoholic fatty liver disease: biochemical, metabolic, and clinical implications. *Hepatology*. 2010;51:679–89.
12. EASL-EASD-EASO. Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol*. 2016;64:1388–402.
13. Farrell GC, Larter CZ. Nonalcoholic fatty liver disease: from steatosis to cirrhosis. *Hepatology*. 2006;43:S99–112.
14. Dyson J, Day C. Treatment of non-alcoholic fatty liver disease. *Digestive Dis*. 2014;32:597–604.
15. Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med*. 2002;346:1221–31.
16. Wong VW-S, Chan RS-M, Wong GL-H, Cheung BH-K, Chu WC-W, Yeung DK-W, et al. Community-based lifestyle modification programme for non-alcoholic fatty liver disease: a randomized controlled trial. *J Hepatol*. 2013;59:536–42.
17. He X-X, Wu X-L, Chen R-P, Chen C, Liu X-G, Wu B-J, et al. Effectiveness of omega-3 polyunsaturated fatty acids in non-alcoholic fatty liver disease: a meta-analysis of randomized controlled trials. *PLoS ONE*. 2016;11:e0162368.
18. Yan J-H, Guan B-J, Gao H-Y, Peng X-E. Omega-3 polyunsaturated fatty acid supplementation and non-alcoholic fatty liver disease: a meta-analysis of randomized controlled trials. *Medicine*. 2018;97:e12271.
19. Parry SA, Hodson L. Influence of dietary macronutrients on liver fat accumulation and metabolism. *J Investigative Med*. 2017;65:1102–15.
20. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med*. 2009;6:e1000100.
21. Green S, Higgins J. Cochrane handbook for systematic reviews of interventions. Version, 2005.
22. ter Horst K, Serlie M. Fructose consumption, lipogenesis, and non-alcoholic fatty liver disease. *Nutrients*. 2017;9:981.
23. Chiu S, Sievenpiper J, De Souza R, Cozma A, Mirrahimi A, Carleton A, et al. Effect of fructose on markers of non-alcoholic fatty liver disease (NAFLD): a systematic review and meta-analysis of controlled feeding trials. *Eur J Clin Nutr*. 2014;68:416.
24. Chung M, Ma J, Patel K, Berger S, Lau J, Lichtenstein AH. Fructose, high-fructose corn syrup, sucrose, and nonalcoholic fatty liver disease or indexes of liver health: a systematic review and meta-analysis. *Am J Clin Nutr*. 2014;100:833–49.
25. Lu W, Li S, Li J, Wang J, Zhang R, Zhou Y, et al. Effects of omega-3 fatty acid in nonalcoholic fatty liver disease: a meta-analysis. *Gastroenterol Res Pract*. 2016;2016.
26. Yu L, Yuan M, Wang L. The effect of omega-3 unsaturated fatty acids on non-alcoholic fatty liver disease: a systematic review and meta-analysis of RCTs. *Pak J Med Sci*. 2017;33:1022.
27. Schrauwen P, van Marken Lichtenbelt W, Saris W, Westerterp KR. Changes in fat oxidation in response to a high-fat diet. *Am J Clin Nutr*. 1997;66:276–82.
28. Schwenzer NF, Springer F, Schraml C, Stefan N, Machann J, Schick F. Non-invasive assessment and quantification of liver steatosis by ultrasound, computed tomography and magnetic resonance. *J Hepatol*. 2009;51:433–45.
29. Reeder SB, Crute I, Hamilton G, Sirlin CB. Quantitative assessment of liver fat with magnetic resonance imaging and spectroscopy. *J Magn Reson Imaging*. 2011;34:729–49.
30. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan —a web and mobile app for systematic reviews. *Syst Rev*. 2016;5:210.
31. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clin trials*. 1986;7:177–88.
32. Luukkonen PK, S.,devirta S, Zhou Y, Kayser B, Ali A, Ahonen L. Saturated fat is more metabolically harmful for the human liver than unsaturated fat or simple sugars. *Diabetes Care*. 2018;41:1732–9.
33. Cohen J. Statistical power analysis for the behavioral sciences. New York, US, Routledge; 2013.
34. Björmo H, Iggman D, Kullberg J, Dahlman I, Johansson L, Persson L, et al. Effects of n-6 PUFAs compared with SFAs on liver fat, lipoproteins, and inflammation in abdominal obesity: a randomized controlled trial. *Am J Clin Nutr*. 2012;95:1003–12.
35. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ: British Medical Journal*. 2003;327:557.
36. Higgins J, Green S. Cochrane handbook for systematic reviews of interventions. Version 5.1.0 [updated March 2011]. The Cochrane Collaboration; 2011. <http://www.cochrane-handbook.org>.
37. Ras RT, Hiemstra H, Lin Y, Vermeer MA, Duchateau GS, Trautwein EA. Consumption of plant sterol-enriched foods and effects on plasma plant sterol concentrations—a meta-analysis of randomized controlled studies. *Atherosclerosis*. 2013;230:336–46.
38. Errazuriz I, Dube S, Slama M, Visentin R, Nayar S, O'connor H, et al. Randomized controlled trial of a MUFA or fiber-rich diet on hepatic fat in prediabetes. *J Clin Endocrinol Metab*. 2017;102:1765–74.
39. Haufe S, Engeli S, Kast P, Böhnke J, Utz W, Haas V, et al. Randomized comparison of reduced fat and reduced carbohydrate hypocaloric diets on intrahepatic fat in overweight and obese human subjects. *Hepatology*. 2011;53:1504–14.
40. Sterne JA, Egger M, Smith GD. Investigating and dealing with publication and other biases. *Systematic Reviews in Health Care: Meta-Analysis in Context*, Second Edition. 2001;323:101–5.
41. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ: British Medical Journal*. 1997;315:629–34.
42. van Herpen NA, Schrauwen-Hinderling VB, Schraut G, Mensink RP, Schrauwen P. Three weeks on a high-fat diet increases intrahepatic lipid accumulation and decreases metabolic flexibility in healthy overweight men. *J Clin Endocrinol Metab*. 2011;96:E691–5.
43. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ: British Medical Journal*. 2011;343:d4002.
44. Bawden S, Stephenson M, Falcone Y, Lingaya M, Ciampi E, Hunter K, et al. Increased liver fat and glycogen stores after consumption of high versus low glycaemic index food: a randomized crossover study. *Diabetes Obes Metab*. 2017;19:70–7.
45. Bendsen NT, Chabanova E, Thomsen HS, Larsen TM, Newman JW, Stender S, et al. Effect of trans fatty acid intake on abdominal and liver fat deposition and blood lipids: a randomized trial in overweight postmenopausal women. *Nutr Diabetes*. 2011;1:e4.
46. Bozzetto L, Prinster A, Annuzzi G, Costagliola L, Mangione A, Vitelli A, et al. Liver fat is reduced by an isoenergetic MUFA diet in a controlled randomized study in type 2 diabetic patients. *Diabetes Care*. 2012;35:1429–35.
47. Kirk E, Reeds DN, Finck BN, Mayurranjan MS, Patterson BW, Klein S. Dietary fat and carbohydrates differentially alter insulin sensitivity during caloric restriction. *Gastroenterology*. 2009;136:1552–60.
48. Marina A, Von Frankenberg AD, Suvag S, Callahan HS, Kratz M, Richards TL, et al. Effects of dietary fat and saturated fat content on liver fat and markers of oxidative stress in overweight/obese men and women under weight-stable conditions. *Nutrients*. 2014;6:4678–90.

49. Markova M, Pivovalova O, Hornemann S, Sucher S, Frahnnow T, Wegner K, et al. Isocaloric diets high in animal or plant protein reduce liver fat and inflammation in individuals with type 2 diabetes. *Gastroenterology*. 2017;152:571–85.e8.

50. Martens EA, Gatta-Cherifi B, Gonnissen HK, Westerterp-Plantenga MS. The potential of a high protein-low carbohydrate diet to preserve intrahepatic triglyceride content in healthy humans. *PloS ONE*. 2014;9:e109617.

51. van Niel M, Feskens EJ, Rietman A, Siebelink E, Mensink M. Partly replacing meat protein with soy protein alters insulin resistance and blood lipids in postmenopausal women with abdominal obesity. *2. J Nutr.* 2014;144:1423–9.

52. Nosaka N, Kasai M, Nakamura M, Takahashi I, Itakura M, Takeuchi H, et al. Effects of dietary medium-chain triacylglycerols on serum lipoproteins and biochemical parameters in healthy men. *Biosci Biotechnol Biochem*. 2002;66:1713–8.

53. Ooi EM, Adams L, Zhu K, Lewis JR, Kerr DA, Meng X, et al. Consumption of a whey protein-enriched diet may prevent hepatic steatosis associated with weight gain in elderly women. *Nutr Metab Cardiovascular Dis*. 2015;25:388–95.

54. Rietman A, Schwarz J, Blokker BA, Siebelink E, Kok FJ, Afman LA, et al. Increasing protein intake modulates lipid metabolism in healthy young men and women consuming a high-fat hypercaloric diet-3. *J Nutr*. 2014;144:1174–80.

55. Rosqvist F, Iggman D, Kullberg J, Cedernaes J, Johansson H-E, Larsson A, et al. Overfeeding polyunsaturated and saturated fat causes distinct effects on liver and visceral fat accumulation in humans. *Diabetes*. 2014;63:2356–68.

56. Utzschneider KM, Bayer-Carter JL, Arbuckle MD, Tidwell JM, Richards TL, Craft S. Beneficial effect of a weight-stable, low-fat/low-saturated fat/low-glycaemic index diet to reduce liver fat in older subjects. *Br J Nutr*. 2013;109:1096–104.

57. Westerbacka J, Lammi K, Häkkinen A-M, Rissanen A, Salminen I, Aro A, et al. Dietary fat content modifies liver fat in overweight nondiabetic subjects. *J Clin Endocrinol Metab*. 2005;90:2804–9.

58. Gepner Y, Shelef I, Komy O, Cohen N, Schwarzfuchs D, Bril N, et al. The beneficial effects of Mediterranean diet over low-fat diet may be mediated by decreasing hepatic fat content. *J Hepatol*. 2019. <https://doi.org/10.1016/j.jhep.2019.04.013>.

59. Schutte S, Esser D, Hoevenaars FPM, Hooiveld GJEJ, Priebe MG, Vonk RJ, et al. A 12-wk whole-grain wheat intervention protects against hepatic fat: the Graandoos study, a randomized trial in overweight subjects. *Am J Clin Nutr*. 2018;108:1264–74. <https://doi.org/10.1093/ajcn/nqy204>.

60. Chen Y, Feng R, Yang X, Dai J, Huang M, Ji X, et al. Yogurt improves insulin resistance and liver fat in obese women with nonalcoholic fatty liver disease and metabolic syndrome: a randomized controlled trial. *Am J Clin Nutr*. 2019;109:1611–9.

61. Marin-Alejandre BA, Abete I, Cantero I, Monreal JI, Elorz M, Herrero JL, et al. The metabolic and hepatic impact of two personalized dietary strategies in subjects with obesity and nonalcoholic fatty liver disease: the fatty liver in obesity (FLiO) randomized controlled trial. *Nutrients*. 2019;11. <https://doi.org/10.3390/nu11102543>.

62. Rosqvist F, Kullberg J, Ståhlman M, Cedernaes J, Heurling K, Johansson H-E, et al. Overeating saturated fat promotes fatty liver and ceramides compared with polyunsaturated fat: a randomized trial. *J Clin Endocrinol Metab*. 2019;104:6207–19.

63. Skytte MJ, Samkani A, Petersen AD, Thomsen MN, Astrup A, Chabanova E, et al. A carbohydrate-reduced high-protein diet improves HbA1c and liver fat content in weight stable participants with type 2 diabetes: a randomised controlled trial. *Diabetologia*. 2019;62:2066–78.

64. Willmann C, Heni M, Linder K, Wagner R, Stefan N, Machann J, et al. Potential effects of reduced red meat compared with increased fiber intake on glucose metabolism and liver fat content: a randomized and controlled dietary intervention study. *Am J Clin Nutr*. 2019;109:288–96. <https://doi.org/10.1093/ajcn/nqy307>.

65. Hodson L, Rosqvist F, Parry SA. The influence of dietary fatty acids on liver fat content and metabolism. *Proc Nutr Soc*. 2019;1–12. <https://doi.org/10.1017/S0029665119000569>.

66. Immamura F, Micha R, Wu JH, de Oliveira Otto MC, Otite FO, Abioye AI, et al. Effects of saturated fat, polyunsaturated fat, monounsaturated fat, and carbohydrate on glucose-insulin homeostasis: a systematic review and meta-analysis of randomised controlled feeding trials. *PLoS Med*. 2016;13:e1002087.

67. Marchesini G, Petta S, Dalle Grave R. Diet, weight loss, and liver health in nonalcoholic fatty liver disease: pathophysiology, evidence, and practice. *Hepatology*. 2016;63:2032–43.

68. Hodson L, Fielding BA. Stearoyl-CoA desaturase: rogue or innocent bystander? *Prog Lipid Res*. 2013;52:15–42.

69. Schwarz JM, Noworolski SM, Wen MJ, Dyachenko A, Prior JL, Weinberg ME, et al. Effect of a high-fructose weight-maintaining diet on lipogenesis and liver fat. *J Clin Endocrinol Metab*. 2015;100:2434–42.